


# Determination of Manganese Content Using a Spectrometer



## Overview

The process involves standardizing potassium permanganate with sodium oxalate and measuring absorbance to create a calibration curve, ultimately quantifying manganese concentration in steel samples. To use spectroscopy to determine the amount of Manganese is an unknown sample. It has a wide range of applications such as wastewater treatment, dye degradation, colloidal nanoparticle characterization. In this application note, a Thermo Scientific™ ARL OPTIMA™ XRF Spectrometer is used to analyze several elements/oxides in ores in a matter of minutes. Manganese (II) reacts with MDTC in sodium acetate-sodium hydroxide buffer of pH 8.0 and gives a brown colored complex.

## Determination of Manganese Content Using a Spectrometer



A simple method is developed to determine trace quantities of Manganese using sodium morpholine dithiocarbamate (Na-MDTC) as a chelating agent by extraction spectrophotometry.



PDF | On Feb 26, 2026, Ajush Dihal published Determination of Manganese Content in Steel Alloy by Spectrophotometric Detection | Find, read and cite all the research you need on...



OPTIM"X XRF Spectrometer with its 13-position sample loader. The first step in manganese production, regardless of fi. al application, is the mining and refinement of manganese ore. Understanding the ...



Explore the spectrophotometric method for determining manganese in steel, including standardization and calibration curve analysis.



In this study, the feasibility of the onsite measurement of nickel, cobalt, manganese, and lithium in an aqueous solution using UV-Vis spectroscopy was investigated for battery material processing and ...



The selectivity of the reaction is studied and the method for determination of manganese (VII) 0.1—2.9 fug/mL is developed. Investigations of extraction in the system discussed were carried ...



Consequently, in this overview we explore the electroanalytical determination of manganese (II) reported throughout the literature and offer insights into future research opportunities ...



The document details a spectrophotometric method for determining manganese in steel, including the preparation of standard solutions, calibration of the spectrophotometer, and construction of ...



In this experiment we will use spectroscopy to measure the amount of manganese in an unknown solution. Schematic diagram of a typical absorption spectrometer.



2.1 Manganese is determined by atomic absorption spectrometry by direct aspiration of the sample into an air-acetylene flame without pre-concentration or pretreatment.

## Contact Us

For more information, pricing, or custom solutions, please contact us:

Website: <https://www.indzawo.co.za>

Email: [sales@indzawo.co.za](mailto:sales@indzawo.co.za)

Phone: +27 71 296 8473

Address: 22 Quantum Street, Midrand, 1685, Gauteng, South Africa

This document is for informational purposes only. Specifications subject to change without notice.

